

# The Effectiveness of Biofloc System Cultivation with the Addition of Supplements Ginger Flour through Feed on the Growth of Tilapia

Cahyono Purbomartono<sup>1\*</sup>, Jejentri<sup>2</sup>, Suwarsito<sup>1</sup>, Dini Siswani Mulia<sup>2</sup>

<sup>1</sup>Department of Aquaculture, Faculty of Agriculture and Fisheries,

<sup>2</sup>Department of Biology Education, Faculty of Education and Teacher,

Universitas Muhammadiyah Purwokerto

Jl. KH. Ahmad Dahlan, PO BOX 202 Purwokerto 53182, Indonesia

\*Corresponding author: [cpurbomartono@yahoo.com](mailto:cpurbomartono@yahoo.com)

## ABSTRACT

DOI;  
[10.30595/jrst.v8i1.19582](https://doi.org/10.30595/jrst.v8i1.19582)

### Histori Artikel:

Diajukan:  
14/10/2023

Diterima:  
25/04/2024

Diterbitkan:  
26/04/2024

*Demand for tilapia consumption cannot be fulfilled due to limited availability. Intensive cultivation is carried out by increasing the stocking density to overcome this. An increase in stocking density can lead to a decrease in water quality which in turn can inhibit growth. To overcome this problem, biofloc system cultivation is carried out with herbal supplements through feed to maintain water quality and optimal fish growth. This study aims to determine the effectiveness of the ginger diet on the growth of tilapia reared for 60 days compared to the ginger and garlic diet for 90 days with biofloc system cultivation. The study was conducted by the experimental method using a completely randomized design, with 3 treatments, 1 control, and each was repeated 4 times for 60 days. The results showed that cultivating tilapia using the biofloc system with the ginger diet for 60 days significantly increased growth, while the garlic diet for 90 days was only significant for weight gain. A ginger diet for 60 days on tilapia and 90 days on gourami with a biofloc cultivation system has the same pattern and effectiveness on growth with an optimal dose of 5.63 g kg<sup>-1</sup> feed. The cultivation of the biofloc system was more effective with the ginger diet for 60 days compared to the ginger and garlic diet for 90 days on fish growth. This shows that biofloc cultivation for 60 days with a ginger diet is recommended for fish farming.*

**Keywords:** Biofloc, Ginger, Garlic, Growth

## 1. INTRODUCTION

Tilapia is a freshwater fish that is widely consumed by the public. The need for tilapia continues to increase with awareness of the consumption of protein derived from fish. Therefore, tilapia cultivation continues to be improved through high stocking densities. High stocking densities are generally followed by additional antibiotics so that growth is maintained. However, antibiotics can cause resistance, environmental pollution, and antibiotic deposits in fish bodies. To overcome this, the use of herbal supplements is now widely used. This is because herbal supplements have advantages over drugs or antibiotics, including being safe to use, and not causing resistance, residue deposits, or environmental pollution.

Several research reports show the use of various herbs for fisheries such as ginger, garlic,

onion, and others. Ginger is a growth promoter rather than an immunostimulant even though it is immunomodulatory (Sukumaran *et al.*, 2016). In principle, other plants are the same and have various bioactivities, but usually, some are more dominant. Research results prove the addition of ginger flour through feed (ginger diet) can increase weight gain, specific growth rate, feed conversion ratio, and increase feed efficiency. Besides growth, a ginger diet strengthens immunity by increasing the total number of leukocytes, lymphocytes, monocytes, and neutrophils. The ginger diet can also increase antioxidant activity which is indicated by increased SOD and CAT (Naliato *et al.*, 2021). Besides ginger, garlic is widely used in the fisheries industry. Garlic is known to increase growth, but garlic is also antiviral, antibacterial, and immunomodulatory (Valenzuela-Gutiérrez

*et al.*, 2021).

Fish cultivators in Indonesia generally have low incomes. Some fish cultivators really hope to benefit from the results of their cultivation. Fast growth, high biomass weight, and survival rate are the hopes of farmers. However, fish and shrimp farming is only sometimes profitable. The alternative is to provide ginger and garlic herbal supplements which are given through feed. To support success, water quality must be maintained properly so that growth can be positive by applying the biofloc cultivation system. The biofloc system is used to improve and maintain good water quality so as to increase feed efficiency and reduce feed conversion. Biofloc converts inorganic nitrogen, especially ammonia, by heterotrophic bacteria into microbial biomass which is then consumed by fish and shrimp as a feed additive (Ekasari, 2009). In addition to using the biofloc system or technology, additional herbal supplements through feed (herbal diet) so that its growth is more optimal.

Previous research has proven that the ginger flour diet (ginger diet) can increase growth by 9.60 g with an optimal dose (DO) of 7.50 g kg<sup>-1</sup> feed for 60 days of study (Purbomartono *et al.*, 2021). As a comparison, for 60 days the research results in garlic supplements only reach weight gain of 8.63 g in tilapia with the optimum dose is 30 g kg<sup>-1</sup> feed for 28 days (Lengka *et al.*, 2013). A similar study using a biofloc system was reported by (Rijal *et al.*, 2021) showed, feeding garlic flour (*Allium sativum*) through feed (garlic diet) with the biofloc system can significantly increase the growth of tilapia (*O. niloticus*) in weight gain of 9.66 g (WG) higher than without biofloc system 4.73 g with optimum dose 40 g kg<sup>-1</sup> feed (Purbomartono *et al.*, 2022). Biofloc is proven to increase the growth and production of tilapia (Sukardi *et al.*, 2018). However, little data shows the combination of treatments to boost growth with a longer maintenance time. The combination of cultivation with the biofloc system and the addition of herbal supplements through feed (diet) for 90 days is expected to produce better growth.

To achieve optimal growth in tilapia aquaculture, a combination of treatments is required. The combination of treatment on the cultivation of the biofloc system with the addition of supplements is still relatively limited. Therefore, this study aims to determine the effectiveness of the ginger diet on the growth of tilapia reared for 60 days compared to the ginger and garlic diet for 90 days with biofloc system cultivation. This research used the dosage of ginger supplements was 75%, 50%, and 25% of

the optimal dose (OD), because the use of a 100% ginger diet has previously been reported to increase significantly the growth of goldfish (Lamin *et al.*, 2018), and tilapia (Payung and Manoppo, 2015). The results of this study are expected to increase the growth of tilapia, maintain water quality, and increase productivity, and the healthier quality of organic fish.

## 2. RESEARCH METHODS

### 2.1 Material

This study used tilapia (7-9 cm), herbal ginger for supplement and the materials for biofloc. Materials for biofloc system consist of probiotic Effective Microorganisms 4 (EM4), molasses and salt. The ginger supplement dose for the treatment based on the results of previous studies, 7.5 g kg<sup>-1</sup> feed (Purbomartono *et al.*, 2021).

### 1.2 Method

Experiments were used as the method of this study, using a completely randomized design with 3 treatments and 1 control, each with 3 replications. This study used 2 combinations, consisting of cultivation with a biofloc system and the addition of ginger flour supplements through feed (ginger diet). The ginger supplements used were 75%, 50% and 25% of the optimal dose (OD), because previously the use of ginger diet (100%) can increase the growth of goldfish (Lamin *et al.*, 2018), and tilapia (Payung and Manoppo, 2015). While biofloc has been shown to increase the growth and production of tilapia (Sukardi *et al.*, 2018).

#### 1.2.1 Experimental design

This study used tilapia which was given the addition of ginger supplements which was given through feed (ginger diet). The full treatment is:

T0: biofloc (control)

T1: biofloc + pellet feed + ginger flour 5.63 (g kg<sup>-1</sup> pakan) (75% OD)

T2: biofloc + pellet feed + ginger flour 3.75 (g kg<sup>-1</sup> pakan) (50% OD)

T3: biofloc + pellet feed + ginger flour 1.88 (g kg<sup>-1</sup> pakan) (25% OD)

#### 2.2.2 Parameter of growth

Measurements of growth consist of weight (WG) gain, length gain (LG) and specific growth rate (SGR) followed Zonneveld *et al.* (1991):

WG = Wt - Wo;

WG = Weight gain

Wt = final weight and

Wo = initial weight

LG = Lt - Lo;

LG = length gain

Lt = final length and

Lo = initial length

SGR =  $[(LnWt-LnWo): t] \times 100\%$

SGR = survival growth rate

t = research time

For feed conversion ratio and feed efficiency:

FE =  $[(Wt-Wo): F] \times 100\%$

FE = feed efficiency

F = total feed consumption

FCR = F:  $[(Wt+D) - Wo]$

FCR = feed conversion rate

### 1.3 Analysis Data

Data were analyzed using Analysis of variance (ANOVA) at a 95% confidence level using the SPSS program. If there is a significant treatment, then proceed with the Duncan Multiple Range Test (DMRT).

## 3. RESULT AND DISCUSSION

### 3.1 Weight gain (WG)

The results of research on the ginger diet with the biofloc system on tilapia for 60 and 90 days and the garlic diet for 90 days resulted in significant WG growth. Observed WG parameter data can be seen in Table 1.

Table 1. Weight gain of tilapia fed ginger diet for 60 days and garlic diet and ginger for 90 days

Treatment	Biofloc system		
	WG Tilapia		WG Gourami**
	Diet ginger 60 days	Diet garlic 90 days*	Diet ginger 90 days
T0	11,058±0,229 <sup>a</sup>	9,57±0,32 <sup>b</sup>	7,19 ± 0,68 <sup>a</sup>
T1	13,440±0,115 <sup>d</sup>	9,13±0,09 <sup>a</sup>	9,42 ± 0,39 <sup>b</sup>
T2	12,758±0,245 <sup>c</sup>	9,14±0,16 <sup>a</sup>	9,02 ± 0,60 <sup>b</sup>
T3	12,533±0,856 <sup>c</sup>	9,66±0,21 <sup>b</sup>	9,20 ± 0,20 <sup>b</sup>

Description: T0 (Control), T1 (pellet feed + ginger flour 5.63 g), T2 (pellet feed + ginger flour 3.75 g), T3 (pellet feed + ginger flour 1.88 g). Different superscripts in the same column shows that there are significant differences (p <0.05). \*Rijal *et al.* (2021); \*\*Purbomartono *et al.* (2022)

Based on Table 1, the ginger diet for 60 days was still better for growth than the garlic diet which was maintained for 90 days with the biofloc system on the growth of tilapia. Even though the fish were different, the 60-day ginger diet (in tilapia) was still better than the 90-day ginger diet (in gourami) for growth which was maintained with the biofloc system. This is a comparison, finding, or preliminary assumption, that giving ginger herbal diet can be done in a shorter time depending on the characteristics of the type of herb and the content of its bioactive compounds. So that it is necessary to develop research for other types of herbs so that they better understand the characteristics of each herb such as dosage, length of time for administration, and method of application.

Ginger contains zingiberene and terpene compounds which produce the characteristic odor and taste of ginger (Jolad *et al.*, 2004). These characteristics of smell and taste can stimulate fish appetite so that feed intake increases. Increased appetite can improve fish growth as indicated by increased weight gain, final weight, relative weight gain, specific growth rate and efficiency, and feed conversion ratio. According to (Stoilova *et al.*, 2007), ginger is known as a

stimulator of digestive enzyme secretion which can increase feed efficiency and growth performance in *Macrobrachium rosenbergii* shrimp (El-Desouky *et al.*, 2012).

The ginger diet of 10-30 g kg<sup>-1</sup> feed in zebrafish was increased the activity of the amylase enzyme (Ahmadifar *et al.*, 2019), indicating that growth performance was responsive to the dose of adding dietary supplements containing ginger. Furthermore, supplementation with 10 g of ginger flour kg<sup>-1</sup> feed in tilapia (El-Sebai *et al.*, 2018) has a positive effect on growth, in trout (*Oncorhynchus mykiss*) at doses of 1-10 g kg<sup>-1</sup> feed, and for white shrimp (*Litopenaeus vannamei*) 5 mg zingerone kg<sup>-1</sup> feed (Chang *et al.*, 2012). Based on the results of this study, the ginger diet was more effective for 60 days than 90 days for tilapia and gourami, the cause cannot be explained with certainty. However, reports from (Naliato *et al.*, 2021), a ginger supplementation diet at the highest levels showed detrimental effects, along with decreased hematological values, and production of H<sub>2</sub>O<sub>2</sub>, NO, SOD, and lysozyme activity. Therefore, it is recommended to administer immunostimulants, in this case ginger powder, considering the dose and duration of

administration. Giving ginger flour immunostimulants through feed with high doses and/or long periods of time can cause immunosuppression which results in decreased growth (Sukumaran *et al.*, 2016).

In contrast to the ginger diet, the garlic diet for 90 days also could not match the growth of tilapia on the ginger diet for 60 days. This is presumably because garlic is more immunomodulatory than growth promoters. As reported by (Foysal *et al.*, 2019), garlic contains bioactive compounds that increase fish immunity against attack by disease agents. In fish farming, garlic is antibacterial, antiparasitic, antioxidant, immunomodulatory, and growth stimulator (Bender-Bojalil and Bárcenas-Pozos, 2013). It

was further explained, to accelerate the immune response, improvement of resistance to certain pathogens can be done by giving garlic extract, raw garlic and certain parts of the ginger plant (Foysal *et al.*, 2019).

### 3.2 Feed Conversion Rate (FCR)

On the FCR indicator, the ginger diet for 60 days and the garlic diet for 90 days had almost the same FCR values. However, the garlic diet for 90 days was not significant to FCR compared to the ginger diet for 60 and 90 days which was significant to FCR. In terms of FCR, it shows that the ginger diet for 90 days is better than 60 days (Table 2).

Table 2. Feed conversion rate of tilapia fed ginger diet for 60 days and garlic diet and ginger for 90 days

Treatment	FCR Tilapia		FCR Gourami**
	Diet ginger 60 days	Diet garlic 90 days*	Diet ginger 90 days
T0	6,55±0,21 <sup>c</sup>	4,22±0,12 <sup>a</sup>	4,82±0,22 <sup>b</sup>
T1	4,38±0,75 <sup>a</sup>	4,36±0,03 <sup>a</sup>	3,87±0,12 <sup>a</sup>
T2	4,87±0,03 <sup>a</sup>	4,37±0,06 <sup>a</sup>	3,98±0,23 <sup>a</sup>
T3	5,30±0,40 <sup>a</sup>	4,29±0,08 <sup>ab</sup>	3,94±0,11 <sup>b</sup>

Description: T0 (Control), T1 (pellet feed + ginger flour 5.63 g), T2 (pellet feed + ginger flour 3.75 g), T3 (pellet feed + ginger flour 1.88 g). Different superscripts in the same column shows that there are significant differences (p <0.05). \*Rijal *et al.* (2021); \*\*Purbomartono *et al.* (2022)

Based on the research results as listed in Table 2, these results raise the notion that FCR is related to the time period of fish rearing. FCR shows the amount of feed needed to increase the weight of 1 kg of fish biomass. To increase the weight of 1 kg of tilapia biomass in this study, a smaller amount of feed was needed in the longer rearing period (90 days) compared to the shorter time (60 days). The results of research conducted by (Jolad *et al.*, 2004), showed that ginger has distinctive smell and taste characteristics that affect the increase in feed intake and feed conversion ratio. Apart from being an attractant (appetite stimulator), ginger is known to increase the activity of digestive enzymes. Described by (Ahmadifar *et al.*, 2019), zebrafish fed a ginger diet of 10-30 g kg<sup>-1</sup> feed could increase the activity of the amylase enzyme, but had no effect on the specific growth rate.

Giving the ginger diet in this study caused the activity of the amylase enzyme which breaks down complex carbohydrate compounds into sugar to increase. The increase in the breakdown of carbohydrates into sugar due to the amylase enzyme causes an increase in the

efficiency of food digestion in the digestive tract of fish. Efficiency in food digestion causes intake and the amount of food needed to decrease in increasing the biomass weight of 1 kg of fish so that the FCR becomes low. This low FCR is more indicative of the performance of digestive enzymes so that the digestibility is better, but does not automatically increase the significance in the growth of tilapia. This is because fish growth is influenced by the quantity and quality of food and the environment. In fish and animals converting organic matter into live biomass, but the effectiveness is influenced by the quality and amount of feed and the environment. The changing environment can affect the type and quality of feed which can affect growth, behavior and physiology of fish (Jobling, 2008).

### 3.3 Feed Efficiency (FE)

Similar to FCR, ginger diet for 60 and 90 days resulted in significant FE but not significant on garlic diet for 90 days (Table 3). Based on the data from this study, feed use efficiency (FE) is more optimal on the 60-day ginger diet compared to the 90-day ginger and garlic diet.

Table 3. Feed Efficiency of tilapia fed ginger diet for 60 days and garlic diet and ginger for 90 days

Treatment	FE Tilapia		FE Gourami**
	Diet ginger 60 days	Diet garlic 90 days*	Diet ginger 90 days
T0	17,78±0,37 <sup>a</sup>	23,73±0,71 <sup>ab</sup>	20,85 ±1,72 <sup>a</sup>
T1	30,63±4,80 <sup>b</sup>	22,94±0,17 <sup>a</sup>	25,84±0,93 <sup>b</sup>
T2	20,46±0,14 <sup>b</sup>	22,87±0,34 <sup>a</sup>	25,16 ± 1,46 <sup>b</sup>
T3	20,10±0,37 <sup>b</sup>	23,30±0,44 <sup>a</sup>	25,41±0,72 <sup>b</sup>

Description: T0 (Control), T1 (pellet feed + ginger flour 5.63 g), T2 (pellet feed + ginger flour 3.75 g), T3 (pellet feed + ginger flour 1.88 g). Different superscripts in the same column shows that there are significant differences (p <0.05). \*Rijal *et al.* (2021); \*\*Purbomartono *et al.* (2022)

Research reports prove that herbs derived from plants can act as a stimulator of digestive enzyme secretion (Stoilova *et al.*, 2007), which affects increased feed efficiency and growth performance in *Macrobrachium rosenbergii* shrimp (El-Desouky *et al.*, 2012). Ginger is an herbal plant that can stimulate the secretion of digestive enzymes in the digestive tract. Apart from being a stimulator of digestive enzymes, ginger is also an attractant which can improve response to food intake, feed efficiency, and better growth performance (Naliato *et al.*, 2021). Reinforced by (Jolad *et al.*, 2004), that dietary ginger supplementation has distinctive odor and taste characteristics which affect the increase in efficiency feed intake and feed conversion ratio, increase final weight, relative weight gain, specific growth rate and protein efficiency rate.

The use of ginger herbal diet at a dose of 0.8% (8 g kg<sup>-1</sup> feed) in koi fish can increase the efficiency of feeding which has an impact on improving growth and production performance. Further explained, the results of research on adding ginger supplements to the feed (ginger diet) can increase the utilization of nutrients as indicated by the increase in body weight, specific growth rate (SGR), protein efficiency ratio (PER), and feed efficiency (FE) (Lamin *et al.*, 2018). In addition to fish, research results on shrimp also show that a combination of ginger *Z. officinalis* and *C. dactylon* supplements can be used as an appetizer, growth booster and immunostimulant so that it effectively increases growth and immunity in *M. rosenbergii* (El-Desouky *et al.*, 2012).

The results of other studies also prove that ginger flour and essential oil can be used as a dietary feed that is anti-inflammatory, antioxidant and growth-promoting (Brum *et al.*, 2017); Sukumaran *et al.*, 2016). Ginger flour supplementation diet in rainbow trout (*Oncorhynchus mykiss*) at a dose of 5 g kg<sup>-1</sup> feed resulted in growth performance and fish resistance to *Aeromonas hydrophila* infection

(Nya and Austin, 2009), while a dose of 10 g kg<sup>-1</sup> feed resulted in significant resistance (Haghighi and Rohani, 2013).

#### 4. CONCLUSION

Cultivating tilapia using the biofloc system with ginger diet for 60 days can significantly increase weight gain, FCR and FE, while the garlic diet for 90 days is only significant for weight gain. A ginger diet for 60 days on tilapia and 90 days on gourami with biofloc cultivation system has the same pattern and effectiveness on growth with optimal dose of 5.63 g kg<sup>-1</sup> feed. The cultivation of the biofloc system was more effective with ginger diet for 60 days compared to the ginger and garlic diet for 90 days on fish growth. Cultivating the biofloc system for 60 days with the ginger diet is recommended for fish farming.

#### ACKNOWLEDGEMENT

I would like to thank to Ms. Aprilliana, Ms. Dhika, Ms. Izanatu, Ms. Shaiffa, Ms. Fitri and Mr. Aan for their technical assistance and research activities. Thanks much also addressed to everyone who helped for success of this research.

#### FUNDING INFORMATION

This research was funded by Universitas Muhammadiyah Purwokerto with grant number: A.11-III/003-S.Pj. /LPPM/I/2021.

#### REFERENCES

- Ahmadifar, E., Sheikhzadeh, N., Roshanaei, K., Dargahi, N., & Faggio, C. (2019). Can dietary ginger (*Zingiber officinale*) alter biochemical and immunological parameters and gene expression related to growth, immunity and antioxidant system in zebrafish (*Danio rerio*). *Aquaculture*, 507 (November 2018), 341-348. <https://doi.org/10.1016/j.aquaculture>.

- 2019.04.049
- Bender-Bojalil, D., & Bárcenas-Pozos, M. E. (2013). El ajo y sus aplicaciones en la conservación de alimentos. *Temas Selectos de Ingeniería de Alimentos*, 7(1), 25–36. Retrieved from <https://docplayer.es/83029-El-ajo-y-sus-aplicaciones-en-la-conservacion-de-alimentos.html>
- Brum, A., Pereira, S. A., Owatari, M. S., Chagas, E. C., Chaves, F. C. M., Mouriño, J. L. P., & Martins, M. L. (2017). Effect of dietary essential oils of clove basil and ginger on Nile tilapia (*Oreochromis niloticus*) following challenge with *Streptococcus agalactiae*. *Aquaculture*, 468, 235–243. <https://doi.org/10.1016/j.aquaculture.2016.10.020>
- Chang, Y. P., Liu, C. H., Wu, C. C., Chiang, C. M., Lian, J. L., & Hsieh, S. L. (2012). Dietary administration of zingerone to enhance growth, non-specific immune response, and resistance to *Vibrio alginolyticus* in Pacific white shrimp (*Litopenaeus vannamei*) juveniles. *Fish and Shellfish Immunology*, 32(2):284–290. <https://doi.org/10.1016/j.fsi.2011.11.017>
- Ekasari, J. (2009). Teknologi Biotlok: Teori dan Aplikasi dalam Perikanan Budidaya Sistem Intensif Bioflocs Technology: Theory and Application in Intensive Aquaculture System. *Jurnal Akuakultur Indonesia*, 8(2), 117–126.
- El-Desouky, H., Abbass, A., El-Asely, A., Gwad, E. A., & Shaheen, A. A. (2012). Effect of *Zingiber officinalis* and *Cyanodon dactylon* against pH stress in giant freshwater prawn *Macrobrachium rosenbergii*. *Proceedings of the 5th Global Fisheries and Aquaculture Research Conference, Faculty of Agriculture, Cairo University, Giza, Egypt*, (5), 1–3.
- El-Desouky, H., El-Asely, A., Shaheen, A. A., & Abbass, A. (2012). Effects of *Zingiber officinalis* and *Cyanodon dactylon* on the Growth Performance and Immune Parameters of *Macrobrachium rosenbergii*. *World Journal of Fish and Marine Sciences*, 4(3), 301–307. <https://doi.org/10.5829/idosi.wjfm.2012.04.03.62120>
- El-Sebai, A., El-Murr, A. E. H., Galal, A. A. A., & Abd El-Motaal, S. M. A. (2018). Effect of ginger dietary supplementation on growth performance, immune response and vaccine efficacy in *Oreochromis niloticus* challenged with *Aeromonas hydrophila*. *Slovenian Veterinary Research*, 55(Suppl 20), 31–39. <https://doi.org/10.26873/SVR-625-2018>
- Foysal, M. J., Alam, M., Momtaz, F., Chaklader, M. R., Siddik, M. A. B., Cole, A., Rahman, M. M. (2019). Dietary supplementation of garlic (*Allium sativum*) modulates gut microbiota and health status of tilapia (*Oreochromis niloticus*) against *Streptococcus iniae* infection. *Aquaculture Research*, 50(8), 2107–2116. <https://doi.org/10.1111/are.14088>
- Haghighi, M., & Rohani, M. S. (2013). The effects of powdered ginger (*Zingiber officinale*) on the haematological and immunological parameters of rainbow trout *Oncorhynchus mykiss*. *J. Med. Plant Herbal Ther. Res*, 1(1), 8–12.
- Jobling, M. (2008). Environmental Factors and Rates of Development and Growth. In *Handbook of Fish Biology and Fisheries*, 1, 97–122. <https://doi.org/10.1002/9780470693803.ch5>
- Jolad, S. D., Lantz, R. C., Solyom, A. M., Chen, G. J., Bates, R. B., & Timmermann, B. N. (2004). Fresh organically grown ginger (*Zingiber officinale*): Composition and effects on LPS-induced PGE2 production. *Phytochemistry*, 65(13), 1937–1954. <https://doi.org/10.1016/j.phytochem.2004.06.008>
- Lamin, B. D., Kumar, M., Pailan, G. H., Singh, D. K., Bisawal, A., & Udit, U. K. (2018). Effect of dietary supplementation of *Zingiber officinale* (Ginger) on growth and nutrient utilization of *Cyprinus carpio* (Linnaeus, 1758). *Journal of Experimental Zoology, India*, 21(2), 849–853.
- Lengka, K., Mannopo, H., & Kolopita, M. E. (2013). Peningkatan Respon Imun Non Spesik Ikan Mas (*Cyprinus carpio* L) Melalui Pemberian Bawang Putih (*Allium Sativum*). *E-Journal Budidaya Perairan*, 1(2), 21–28. <https://doi.org/10.35800/bdp.1.2.2013.1912>
- Naliato, R. F., Carvalho, P. L. P. F., Vicente, I. S. T., Xavier, W. dos S., Guimarães, M. G., Rodrigues, E. J. D., Barros, M. M. (2021).

- Ginger (*Zingiber officinale*) powder improves growth performance and immune response but shows limited antioxidant capacity for Nile tilapia infected with *Aeromonas hydrophila*. *Aquaculture Nutrition*, 27(3), 850–864. <https://doi.org/10.1111/anu.13229>
- Nya, E. J., & Austin, B. (2009). Use of dietary ginger, *Zingiber officinale* Roscoe, as an immunostimulant to control *Aeromonas hydrophila* infections in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Journal of Fish Diseases*, 32(11), 971–977. <https://doi.org/10.1111/j.1365-2761.2009.01101.x>
- Payung, C. N., & Manoppo, H. (2015). Peningkatan Respon Kebal Non-spesifik dan Pertumbuhan Ikan Nila (*Oreochromis niloticus*) Melalui Pemberian Jahe, *Zingiber officinale*. *Jurnal Budidaya Perairan*, 3(1), 11–18.
- Purbomartono, C., Habibah, U., Wahyuningtyas, R. A., Husin, A., & Samadan, G. M. (2021). Growth and immunity of african catfish (*Clarias gariepinus*) with dietary inclusion of ginger (*Zingiber officinalis*) and turmeric (*Curcuma domestica*). *AAFL Bioflux*, 14(3), 1365–1372.
- Purbomartono, C., Hapsari, A. N., Susanto, & Samadan, G. M. (2022). Ginger (*Zingiber officinale* Rosc.) flour in Gourami (*Osporonemus gourami*) hatchery with biofloc System. *AAFL Bioflux*, 15(2), 585–592.
- Purbomartono, C., Husin, A., Bagasnabila, I. S., Zularini, F. G. D., Susiyani, A. T., Purwaningsih, E. P., & Purnomo, P. (2022). Efektivitas dan Potensi Herbal untuk Peningkatan Pertumbuhan Benih Lele dumbo (*Clarias gariepinus*). *Sainteks*, 19(2), 219–229. <https://doi.org/10.30595/sainteks.v19i2.14903>
- Rijal, M. A., Purbomartono, C., & Jannah, I. F. (2021). Respon Pertumbuhan Ikan Nila (*Oreochromis niloticus*) yang diberi Pakan Supplementasi Bawang Putih (*Allium sativum*) pada Sistem Bioflok. *Sainteks*, 18(2), 117–122. <https://doi.org/10.30595/sainteks.v18i2.12773>
- Stoilova, I., Krastanov, A., Stoyanova, A., Denev, P., & Gargova, S. (2007). Antioxidant activity of a ginger extract (*Zingiber officinale*). *Food Chemistry*, 102(3), 764–770. <https://doi.org/10.1016/j.foodchem.2006.06.023>
- Sukardi, P., Soedibya, P. H. T., & Pramono, T. B. (2018). Produksi Budidaya Ikan Nila (*Oreochromis niloticus*) Sistem Bioflok Program Studi Budidaya Perairan Fakultas Perikanan dan Ilmu Kelautan. *Asian Journal of Innovation and Entrepreneurship*, 3(2), 198–203.
- Sukumaran, V., Park, S. C., & Giri, S. S. (2016). Role of dietary ginger *Zingiber officinale* in improving growth performances and immune functions of *Labeo rohita* fingerlings. *Fish and Shellfish Immunology*, 57, 362–370. <https://doi.org/10.1016/j.fsi.2016.08.056>
- Valenzuela-Gutiérrez, R., Lago-Lestón, A., Vargas-Albores, F., Cicala, F., & Martínez-Porchas, M. (2021). Exploring the garlic (*Allium sativum*) properties for fish aquaculture. *Fish Physiology and Biochemistry*, 47(4), 1179–1198. <https://doi.org/10.1007/s10695-021-00952-7>
- Zonneveld N., Huisman E. A., Boon J. H. (1991). *Fish Cultivation Principles*. Jakarta: PT Gramedia Pustaka Utama